

Time-domain astronomy

Sylvia J. Zhu

DESY Zeuthen

sylvia.zhu@desy.de

[she/her]



Rough outline

Day 1: Intro + detectors

How are gamma rays produced and how do we detect them?

Day 2: Rapid transients

How do IACTs observe gamma-ray bursts and other rapidly fading transients?


Day 3: Longer duration transients

What else must be considered for longer duration transients like supernovae?

core collapse supernovae
flaring AGN as gravitational-wave candidates
maybe other stuff???

before I begin

I found some old pictures of other lecturers




David in 2011



David and Fabio in 2012

Part 3a. Core collapse supernovae

How should we observe slowly evolving transients?



From novae to supernovae?

where are the gamma rays

So, we can get gamma rays from novae

What about supernovae?

nova energy release $\sim 10^{42}$ erg

supernova energy release $\sim 10^{51}$ erg

side note


“foe” = (10 to the) fifty-one erg

nova distance \sim kpc

supernova distance \sim Mpc


=> from energy arguments, a detection is plausible!

So, are the right ingredients there?



Core-collapse supernovae from astronomy 101


star goes boom




Core-collapse supernovae from astronomy 101

star goes boom

Massive stars fuses successively heavier elements until iron



R. J. Hall




© 2012 Encyclopædia Britannica, Inc.

Core-collapse supernovae from astronomy 101


star goes boom

Massive stars fuses successively heavier elements until iron

R. J. Hall



R. J. Hall



stuff happens*, ultimately a lot of material is ejected at relativistic speeds

*complex, extreme physical conditions, difficult to properly model

Core-collapse supernovae from astronomy 101

star goes boom



modified from J. Borowska-Naguszewska


Massive star explodes as a supernova, lightcurve is bright at the start and then simply fades?

Core-collapse supernovae from astronomy 101


star goes boom

102

Any non-Ia supernova is a core collapse supernova



R. J. Hall




A. Villar

Core-collapse supernovae from astronomy 101


star goes boom

102


Any non-Ia supernova is a core collapse supernova



R. J. Hall



A. Villar




Core-collapse supernovae from astronomy 101


star goes boom

102

Any non-Ia supernova is a core collapse supernova



R. J. Hall



stripped
envelope
supernovae


A. Villar

Core-collapse supernovae from astronomy 101


star goes boom

102

Any non-Ia supernova is a core collapse supernova



R. J. Hall





A. Villar

Core-collapse supernovae from astronomy 101

star goes boom

102



Supernovae are brightest at ~optical wavelengths

Emission is powered by **radioactive decay**

neutron capture ->
increase mass number A

beta decay to a more stable element ->
increase atomic number Z
+
nucleus drops to lower energy state ->
release photons

[M. Modjaz et al., Nature Astronomy 3 (2019)]

Pause to talk about nucleosynthesis

How do we get elements heavier than helium?

Group	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	
1	1 H																		2 He
2	3 Li	4 Be											5 B	6 C	7 N	8 O	9 F	10 Ne	
3	11 Na	12 Mg											13 Al	14 Si	15 P	16 S	17 Cl	18 Ar	
4	19 K	20 Ca	21 Sc	22 Ti	23 V	24 Cr	25 Mn	26 Fe	27 Co	28 Ni	29 Cu	30 Zn	31 Ga	32 Ge	33 As	34 Se	35 Br	36 Kr	
5	37 Rb	38 Sr	39 Y	40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	50 Sn	51 Sb	52 Te	53 I	54 Xe	
6	55 Cs	56 Ba	* 71 Lu	72 Hf	73 Ta	74 W	75 Re	76 Os	77 Ir	78 Pt	79 Au	80 Hg	81 Tl	82 Pb	83 Bi	84 Po	85 At	86 Rn	
7	87 Fr	88 Ra	* 103 Lr	104 Rf	105 Db	106 Sg	107 Bh	108 Hs	109 Mt	110 Ds	111 Rg	112 Cn	113 Nh	114 Fl	115 Mc	116 Lv	117 Ts	118 Og	
			* 57 La	58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb			
			* 89 Ac	90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No			

[Double sharp]

Pause to talk about nucleosynthesis

How do we get elements heavier than helium?


Group	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
Period 1	1 H	formed in the Big Bang (“Big Bang nucleosynthesis”)																2 He
2	3 Li	4 Be											5 B	6 C	7 N	8 O	9 F	10 Ne
3	11 Na	12 Mg											13 Al	14 Si	15 P	16 S	17 Cl	18 Ar
4	19 K	20 Ca	21 Sc	22 Ti	23 V	24 Cr	25 Mn	26 Fe	27 Co	28 Ni	29 Cu	30 Zn	31 Ga	32 Ge	33 As	34 Se	35 Br	36 Kr
5	37 Rb	38 Sr	39 Y	40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	50 Sn	51 Sb	52 Te	53 I	54 Xe
6	55 Cs	56 Ba	* 71 Lu	72 Hf	73 Ta	74 W	75 Re	76 Os	77 Ir	78 Pt	79 Au	80 Hg	81 Tl	82 Pb	83 Bi	84 Po	85 At	86 Rn
7	87 Fr	88 Ra	* 103 Lr	104 Rf	105 Db	106 Sg	107 Bh	108 Hs	109 Mt	110 Ds	111 Rg	112 Cn	113 Nh	114 Fl	115 Mc	116 Lv	117 Ts	118 Og
			* 57 La	58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb		
			* 89 Ac	90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No		

????

[Double sharp]

Pause to talk about nucleosynthesis


How do we get elements heavier than helium?



Pause to talk about nucleosynthesis

How do we get elements heavier than helium?

key ingredient: free neutrons are captured by nuclei




[P. Möller, from J. Cowan and F.-H. Thielemann, Physics Today 57 (2004)]

Pause to talk about nucleosynthesis

How do we get elements heavier than helium?


An example of an r-process nucleosynthesis simulation



[J. Lippuner, SkyNet]

Pause to talk about nucleosynthesis

How do we get elements heavier than helium?




Core-collapse supernovae from astronomy 101


star goes boom

102

Any non-Ia supernova is a core collapse supernova



R. J. Hall




A. Villar

Core-collapse supernovae from astronomy 101


star goes boom

102


Any non-Ia supernova is a core collapse supernova



R. J. Hall



A. Villar




why are some SNe so bright / long lasting?
are additional *interactions* providing extra power?

Core-collapse supernovae from astronomy 101


star goes boom

102


Any non-Ia supernova is a core collapse supernova



R. J. Hall



A. Villar




narrow lines indicate slow moving gas:
atoms are ionized by SN ejecta, lose an electron
ions recapture electrons (recombination)
electron cascades down into lower energy levels
-> emission at specific wavelengths

Core-collapse supernovae from astronomy 101


star goes boom

102

Any non-Ia supernova is a core collapse supernova



R. J. Hall



A. Villar

some supernovae show evidence for additional interactions with (slow-moving) external material
=> interacting supernovae

Core-collapse supernovae from astronomy 101

star goes boom




modified from J. Borowska-Naguszewska

Massive star explodes as a supernova, lightcurve is bright at the start and then simply fades?

Core-collapse supernovae from astronomy 101

star goes boom

Massive stars emit strong stellar winds over their lifetimes



line driven winds: photons are absorbed and the average momentum gain is outward



modi

Massive

These winds create densities in the circumstellar environment


Core-collapse supernovae from astronomy 101

star goes boom



Core-collapse supernovae from astronomy 101

star goes boom




modified from J. Borowska-Naguszewska

Expanding SN ejecta runs into circumstellar material -> shocks -> particle acceleration -> gamma rays????

Core-collapse supernovae from astronomy ~~101~~¹⁰²

star goes boom



- Gamma rays have been detected from novae
- Supernovae should have dense material in their surrounding environment -> shocks

So, have we detected gamma rays from supernovae?


modified from J. Borowska-Naguszewska

Expanding SN ejecta runs into circumstellar material -> shocks -> particle acceleration -> gamma rays????

Searches for gamma-ray emission from CC SNe


Fermi-LAT observations of SN 2023ixf

A SN II-n just 7 Mpc away ($z = 0.0008$)



[K. Murase et al., MNRAS 440 (2014)]

strong emission lines that faded over time



[N. Smith et al., ApJ 956 (2023)]



Searches for gamma-ray emission from CC SNe

Fermi-LAT observations of SN 2023ixf

LAT upper limits on >100 MeV flux \rightarrow upper limits on efficiency of cosmic ray acceleration

how much of the energy goes into accelerating cosmic rays

[G. Martí-Devesa, A&A 686 (2024)]




Searches for gamma-ray emission from CC SNe

Fermi-LAT observations of SN 2023ixf

LAT upper limits on >100 MeV flux \rightarrow upper limits on efficiency of cosmic ray acceleration of $\lesssim 1\%$



[G. Martí-Devesa, A&A 686 (2024)]



Complication: γ - γ opacity

When should we look for VHE gamma rays?

The supernova itself is bright in optical-ish photons \rightarrow γ rays are absorbed initially (~10s to 100s of days?)




rule of thumb for γ - γ pair production:
 $E_1 E_2 \gtrsim (0.5 \text{ MeV})^2$

Complication: γ - γ opacity


When should we look for VHE gamma rays?

The supernova itself is bright in optical-ish photons \rightarrow γ rays are absorbed initially (~10s to 100s of days?)

e.g., the recent RS Ophiuchi nova event



[H.E.S.S. Collaboration, Science 376 (2022)]




rule of thumb for γ - γ pair production:

$$E_1 E_2 \gtrsim (0.5 \text{ MeV})^2$$

Complication: γ - γ opacity

When should we look for VHE gamma rays?

The supernova itself is bright in optical-ish photons \rightarrow γ rays are absorbed initially (~ 10 s to 100s of days?)




Actual predictions depend strongly on optical SN properties, circumstellar environment, etc.

Searches for gamma-ray emission from CC SNe

H.E.S.S. observations of CC SNe

VHE photons from CC SNe need to wait until the opacity decreases sufficiently (~10s to 100s of days?)

H.E.S.S. observations:




[H.E.S.S. Collaboration, A&A 626 (2019)]

Searches for gamma-ray emission from CC SNe

H.E.S.S. observations of CC SNe

VHE photons from CC SNe need to wait until the opacity decreases sufficiently (~10s to 100s of days?)

H.E.S.S. upper limits on mass loss rate for an efficiency of 10% (+ other assumptions)




[H.E.S.S. Collaboration, A&A 626 (2019)]

Searches for gamma-ray emission from CC SNe

MAGIC observations of SN 2014J

VHE photons from CC SNe need to wait until the opacity decreases sufficiently (~10s to 100s of days?)

MAGIC observed a very nearby SN Ia (3.6 Mpc):




ally (Dwarkadas & Chevalier 1998). Although the flux keeps increasing with time according to Eq. (1), with this low expansion parameter it will still be about 10^{-21} photons $\text{cm}^{-2} \text{s}^{-1}$ 100 yr after the SN occurred, which is well below the sensitivity of the current and planned VHE observatories – several orders of magnitude below the sensitivity that the future Cherenkov Telescope Array (CTA¹) will reach.

No detection -> no VHE detection possible for any SN Ia-CSM?

[M. L. Ahnen et al., A&A 602 (2017)]

Circumstellar density profile is complex

Shells from stellar winds of massive stars




- Wind mass loss is episodic, not continuous
-> “shells”
- Shape is affected by environment (e.g., stellar companions)

[L. Decin, ESO, ALMA]

Circumstellar density profile is complex

Interactions with circumstellar shells -> late-time rebrightening

e.g., optical images of SN1987A





SNe can rebrighten after months or years, when the SN shock interacts with a dense circumstellar shell

NASA, ESA, R. Kirshner (CfA, Gordon and Betty Moore Foundation), P. Challis (CfA)

Prospects for gamma rays from rebrightening supernovae



Interactions with circumstellar shells -> late-time rebrightening



[R. Brose et al., A&A 706 (2026)]

Prospects for gamma rays from rebrightening supernovae

Interactions with circumstellar shells -> late-time rebrightening



[R. Brose et al., A&A 706 (2026)]


Depending on the shell properties, this could also produce detectable gamma-ray emission



We are starting to explore different observation strategies for SNe (e.g., observe later and/or observe based on MWL rebrightening)

Prospects for gamma rays from rebrightening supernovae

Interactions with circumstellar shells -> late-time rebrightening



Some core-collapse supernovae have **rebrightened** after ~years

Rebrightened flux can even be much higher than original SN flux (in radio)

-> interactions with circumstellar shells

“late-time supernova rebrightening”










≈

“very young supernova remnant”


Prospects for gamma rays from rebrightening supernovae

Interactions with circumstellar shells -> late-time rebrightening

SN 2007it on the RISE - a radio detection of an interacting supernova 18 years post-explosion

F. ACERO ¹ R. Z. E. ALSABERI ^{2,3} M. ARIAS ⁴ J. BOROWSKA-NAGUSZEWSKA,⁵ R. BROSE ^{6,*}
C. BURGER-SCHIEDLIN ^{7,8} P. G. EDWARDS ⁹ Q. FENG ¹⁰ M. D. FILIPOVIĆ ² T. LASKAR ¹⁰
S. LAZAREVIĆ ^{2,9,11,†} J. MACKEY ^{7,8} A. NUCARA ^{12,13} K. ROSE ^{14,9,‡} S. RYDER ^{15,16} F. SCHÜSSLER ¹⁷
A. SIMONGINI ^{18,13} Z. J. SMEATON ² I. SUSHCH ^{19,20,21} AND S. ZHU ²²

THE RISE COLLABORATION




[F. Acero et al., RNAAS 10 (2026)]

Core-collapse supernovae from astronomy ~~101~~

star goes boom

102



modified from J. Borowska-Naguszowska

Usually the answer is “No” and I end my SN slides here ...

Expanding SN ejecta

... but WAIT, Fabio has some exciting news going public today ...


-> gamma rays????

Core-collapse supernovae from astronomy 101


star goes boom

102


Any non-Ia supernova is a core collapse supernova



R. J. Hall




A. Villar



why are some SNe so bright / long lasting?
are additional *interactions* providing extra power?

What powers a superluminous supernova?

go ask Fabio




[A. Gal-Yam, Science 337 (2012)]

- circumstellar shell interactions?
- accretion onto the new-born central compact object?
- magnetar central engine?


Have we detected gamma rays from a superluminous supernova?

go ask Fabio

the host galaxy normally looks like this



the supernova is clearly visible in optical light



gamma-ray excess 🙄


Prospects for CTAO

these slides are by Fabio, ask him for more info

What's the best way to observe a supernova?


Significance for each night

5 hours every night. Analysis between 30 GeV - 100 TeV



13


Deeper observations : 40h over 5 nights



15

Horizon of detectability

40 hours total time. Starting from $T=20$ days \Rightarrow 28d




Horizon of 7 Mpc for Type IIp and 70 Mpc for Type IIIn

- Probe the starburst galaxies M82, NGC 253, NGC 6946 @ < 7 Mpc
- For cosmological scales this means :
 - Probing the local sheet (< 10 Mpc) & core of Laniakea at 70 Mpc


Core-collapse supernovae from astronomy 101

102

Any non-Ia supernova is a core collapse supernova




R. J. Hall




A. Villar

Core-collapse supernovae from astronomy 101 ~~102~~ 999

Any non-Ia supernova is a core collapse supernova




R. J. Hall




A. Villar

a large fraction of stars are in binaries




Z.-W. Han et al., Res Astron Astrophys 20 (2020)

binary evolution



binary evolution



If a large fraction of stars are in binaries, then a large fraction of SNe should occur in binaries => major differences in star's evolution, environment, properties at explosion, etc.

- AGB: asymptotic giant branch
- B: helium star
- BH: black hole
- BS: blue straggler
- CE: common envelope
- CEEMP: carbon-enhanced metal-poor star
- CV: cataclysmic variable
- DCO: double compact objects
- FRB: fast radio burst
- GRB: gamma-ray burst
- GWR: gravitational wave radiation
- He/CO-WD: helium or carbon oxygen white dwarf
- He-Star: helium star
- HM: Hertzsprung gap
- HMXB: high mass X-ray binary
- L/IMXB: low- or intermediate mass X-ray binary
- MS: main sequence
- NS: neutron star
- RGB: red giant branch
- RLOF: Roche-lobe over flow
- RSG: red supergiant
- sdOB: hot subdwarf O/B
- SG: supergiant
- SN: supernova
- Symb: symbiotic star
- TZD: Thorne-Zytkow object
- WD: white dwarf
- WMT: wind mass transfer

$q = M_{\text{donor}}/M_{\text{recipient}}$ at the onset of mass transfer.

For MS/SG massive stars, q , varies from less than 20. For RGB massive stars, q , varies from 1.4 to 3.

For MS/SG low- and intermediate-mass stars ($M < 0.6M_{\odot}$), if $M < 1.6M_{\odot}$, q , varies from 2 to 5; if $M < 1.6M_{\odot}$, q , varies from 1 to 4.


For RGB/AGB low- and intermediate-mass stars, q , varies from 1 to 3.

1. There is a possibility that pair instability in massive stars leaves no remnant.
2. There is a possibility that the binary system is disrupted by the supernova explosion and its kick.
3. There is a possibility that He-Star RLOF may occur, but we omit this in the diagram for simplicity.


Supernovae in binaries are important

Time-domain astronomy in the era of GW detections

How do the gravitational-wave binaries form and evolve?



LIGO-Virgo-KAGRA / Aaron Geller / Northwestern




M. U. Kruckow et al., MNRAS 481 (2018)

The curious case of SN2022jli

CC SNe in a binary system


P. Chen et al., Nature 625 (2024)



The curious case of SN2022jli

P. Chen et al., Nature 625 (2024)

CC SNe in a binary system




The curious case of SN2022jli

P. Chen et al., Nature 625 (2024)

CC SNe in a binary system

Interpretation: Compact object remnant is accreting from companion in a binary w/ a 12.4-day orbit




ESO / L. Calçada

The curious case of SN2022jli

CC SNe in a binary system

P. Chen et al., Nature 625 (2024)

Hint of gamma rays six months after explosion?



Interpretation: jet or shocks
-> particle acceleration
-> gamma rays (once γ - γ opacity has decreased sufficiently)




Is this another channel for gamma-ray emission from core collapse SNe?

Pause: return to GRBs

When should we look for VHE gamma rays?

γ - γ opacity is an issue whenever there are a lot of ambient photons at the “right” energy

q: what about for GRBs?




[NASA's Goddard Space Flight Center]

Indeed, the GRB jet is initially **optically thick**, photons only escape once jet has expanded and density has decreased sufficiently

-> for GRBs, the timescale for sufficiently reduced γ - γ opacity is $< s$

Pause: return to GRBs


How are long GRBs and core-collapse supernovae related?



A. Villar

The SNe associated with long GRBs tend to be missing both hydrogen and helium lines (“stripped envelope” supernovae)

“BL” = broad lines -> very fast moving material



Pause: return to GRBs

How are long GRBs and core-collapse supernovae related?



[DESY, Science Communication Lab]


Pause: return to GRBs

How are long GRBs and core-collapse supernovae related?


Most GRB-SN are ordinary or even subluminal

Exception: Are superluminous supernovae connected with ultralong GRBs?

“ultralong”: duration ~ 1000 s of seconds



[G. Stratta et al., ApJ 779 (2013)]



[D. A. Kann, A&A 624 (2019)]



Pause: return to GRBs

How are long GRBs and core-collapse supernovae related?

Most GRB-SN are

Exception: Are s

“ultralong”: d



[D. A. Kann, A&A 624 (2019)]

Days after GRB Trigger in the Rest-Frame



[D. A. Kann, A&A 624 (2019)]



[G. Stratta et al., ApJ 779 (2013)]

Pause: return to GRBs


How are long GRBs and core-collapse supernovae related?

Most GRB-SN are


Exception: Are s
“ultralong”: d



are magnetars the key?



[ESO/L. Calçada]



[D. A. Kann, A&A 624 (2019)]

[D. A. Kann, A&A 624 (2019)]

[G. Stratta et al., ApJ 779 (2013)]

Part 3b. Flaring AGN as gravitational-wave sources




Another messenger related to AGN (maybe)

ok here is my disclaimer that I know very little about AGN


go ask Lea

In a nutshell: (super)massive black hole in the middle of a galaxy accretes material and emits photons




[Emma Alexander]

How we name it depends on the system's properties **and** on our viewing angle



[ESO/WFI; MPIfR/ESO/APEX/A.Weiss et al.; NASA/CXC/CfA/R.Kraft et al.]



[ESA/Hubble & NASA]



AGN are also major time domain astronomy sources

go ask Lea

AGN often exhibit multiwavelength flaring activity

all the IACTs have active programs for observing AGN (mostly blazars) during flares


- jet precession?
- stochastic processes inside the jet?
- evidence of a binary companion??



[NASA]

AGN with (quasi)periodic oscillations could become GW sources (super)massive black hole binaries


OJ 287 is a blazar with quasiperiodic oscillations (~12 year period?)




[A. Sillinpää et al., ApJ 325 (1988)]

AGN with (quasi)periodic oscillations could become GW sources (super)massive black hole binaries

OJ 287 is a blazar with quasiperiodic oscillations (~12 year period plus 60-year envelope?)





[A. Sillinpää et al., ApJ 325 (1988)]



[L. Dey et al., ApJ 866 (2018)]

AGN with (quasi)periodic oscillations could become GW sources (super)massive black hole binaries

OJ 287 is a blazar with quasiperiodic oscillations (~12 year period plus 60-year envelope?)





[L. Dey et al., ApJ 866 (2018)]

AGN with (quasi)periodic oscillations could become GW sources


(super)massive black hole binaries

OJ 287 is a blazar with quasiperiodic oscillations (~12 year period plus 60-year envelope?)



AGN with (quasi)periodic oscillations could become GW sources (super)massive black hole binaries

A few other candidates have also been found, e.g. J1048+7143




[E. Kun et al., ApJ 940 (2022)]

[E. Kun et al., ApJL 963 (2024)]

AGN with (quasi)periodic oscillations could become GW sources (super)massive black hole binaries

A few other candidates have also been found, e.g. J1048+7143



[E. Kun et al., ApJ 940 (2022)]

[E. Kun et al., ApJL 963 (2024)]

flare was predicted here


source is indeed flaring right now 🗨️

to learn more: talk to Ilja Jaroschewski
(CEA Saclay)

GWs from (S)MBHBs

Scaling up the LIGO/Virgo/KAGRA black hole mergers

Reminder: Anything with a changing mass quadrupole moment produces GWs



Animation created by SXS, the Simulating eXtreme Spacetimes (SXS) project (<http://www.black-holes.org>)
Video and explanation: <https://www.ligo.caltech.edu/video/ligo20160211v10>


THE SPECTRUM OF GRAVITATIONAL WAVES

Observatories & experiments

Ground-based experiment



Space-based observatory



Pulsar timing array



Cosmic microwave background polarisation



Timescales

milliseconds

seconds

hours

years

billions of years

Frequency (Hz)

100

1

10^{-2}

10^{-4}


10^{-6}

10^{-8}

10^{-16}

Cosmic fluctuations in the early Universe

Cosmic sources




Supernova



Pulsar



Compact object falling onto a supermassive black hole



Merging supermassive black holes



Merging neutron stars in other galaxies



Merging stellar-mass black holes in other galaxies



Merging white dwarfs in our Galaxy

AGN w ~periodic variability?


LISA gravitational-wave observatory

Laser Interferometer Space Antenna



ground-based (LIGO/Virgo/KAGRA)

effective arms: $O(1000 \text{ km})$
 $\lambda \sim 1000 \text{ km} \rightarrow f_{\text{GW}} \sim O(100 \text{ Hz})$




space-based (LISA)

effective arms: $O(2.5e6 \text{ km})$
 $\lambda \sim 10^6 \text{ km} \rightarrow f_{\text{GW}} \sim O(0.1 \text{ Hz})$

LISA gravitational-wave observatory

Laser Interferometer Space Antenna



gro
effe
 $\lambda \sim$

finally: some other things


obviously time-domain astronomy in gamma rays is much more than what we've discussed here

here is a parade of some of those other things

T D E

tidal disruption events (TDEs)

another way to potentially get gamma rays from supermassive black holes




[S. Komossa, JHEAP 7 (2015)]

tidal disruption events (TDEs)

another way to potentially get gamma rays from supermassive black holes

Observing programs are underway for all IACTs
BUT gamma-gamma opacity is again an issue -> when to observe?



A few examples of IACT searches:
MAGIC collaboration, A&A 552 (2013)
I. Lypova et al., Gamma2022


[S. Komossa, JHEAP 7 (2015)]

tidal disruption events (TDEs)

side note on TDEs as potential neutrino sources

Are (some) IceCube-detected neutrinos produced by TDEs? We used to think so, now we don't ...

[A. Zegarelli et al., ICRC2025]



In the meantime, the IceCube collaboration improved their localization algorithms ...


fast radio bursts (FRBs)

we basically have no idea what they are*

* except for the one associated with a soft gamma repeater

a magnetar that produces a burst of (soft) gamma rays

a neutron star with a very high magnetic field



- very short (<seconds) bursts of radio
- seem to be at cosmological distances
- some fraction of FRB sources are *repeaters* (multiple bursts)

see this review by J. I. Katz for more info

[D. R. Lorimer, Nature Astronomy 2 (2018)]

fast radio bursts (FRBs)


we basically have no idea what they are*

* except for the one associated with a soft gamma repeater

IACTs mostly have to look for some sort of “afterglow” emission

exception: VERITAS can observe ~simultaneously with CHIME

nothing so far 🙄



- very short (<seconds) bursts of radio
- seem to be at cosmological distances
- some fraction of FRB sources are *repeaters* (multiple bursts)

see [this review](#) by J. I. Katz for more info

[D. R. Lorimer, Nature Astronomy 2 (2018)]